

## Key Problems

### Problem 1.

Find all eigenvalues of  $A$ , and a base for the eigenspace  $E_\lambda$  for each eigenvalue  $\lambda$ :

$$\text{a) } A = \begin{pmatrix} 5 & 9 \\ 9 & 5 \end{pmatrix}$$

$$\text{b) } A = \begin{pmatrix} 2 & 1 \\ -1 & 4 \end{pmatrix}$$

$$\text{c) } A = \begin{pmatrix} 3 & -4 \\ 3 & 0 \end{pmatrix}$$

$$\text{d) } A = \begin{pmatrix} 3 & 0 & 1 \\ 0 & 4 & 0 \\ 1 & 0 & 3 \end{pmatrix}$$

$$\text{e) } A = \begin{pmatrix} 0 & 1 & 1 \\ 1 & 0 & 1 \\ 1 & 1 & 0 \end{pmatrix}$$

$$\text{f) } A = \begin{pmatrix} 0 & 1 & 1 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix}$$

### Problem 2.

For the matrix  $A$  in Problem 1 a) - f), determine whether  $A$  is diagonalizable, and find an invertible matrix  $P$  and a diagonal matrix  $D$  such that  $P^{-1}AP = D$  when this is possible.

### Problem 3.

Find the eigenvalues of  $A$ , and show that  $A$  is diagonalizable:

$$A = \begin{pmatrix} 1 & 0 & 0 & 4 \\ 0 & 2 & 3 & 0 \\ 0 & 3 & 2 & 0 \\ 4 & 0 & 0 & 1 \end{pmatrix}$$

### Problem 4.

Use eigenvalues and eigenvectors of  $A$  to determine the limit of  $A^m$  when  $m \rightarrow \infty$ .

$$\text{a) } A = \begin{pmatrix} 0.40 & 0.15 \\ 0.60 & 0.85 \end{pmatrix}$$

$$\text{b) } A = \begin{pmatrix} 0.77 & 0.46 \\ 0.23 & 0.54 \end{pmatrix}$$

### Problem 5.

Show that when  $A$  is a  $3 \times 3$  matrix, then the characteristic equation of  $A$  can be written as  $-\lambda^3 + c_1\lambda^2 - c_2\lambda + c_3 = 0$ , where  $c_1 = \text{tr}(A)$ ,  $c_2 = M_{12,12} + M_{23,23} + M_{13,13}$  and  $c_3 = \det(A)$ . Hint: Write down the characteristic equation of a  $3 \times 3$  matrix  $A = (a_{ij})$  with general coefficients. Then use the formula to find the characteristic equation and the eigenvalues of the following matrices:

$$\text{a) } A = \begin{pmatrix} 1 & 1 & 1 \\ 2 & 4 & 0 \\ 3 & 5 & 1 \end{pmatrix}$$

$$\text{b) } A = \begin{pmatrix} 3 & 2 & 1 \\ 2 & 3 & 2 \\ 1 & 2 & 3 \end{pmatrix}$$

$$\text{c) } A = \begin{pmatrix} 0 & 4 & 7 \\ 0 & 0 & 5 \\ 0 & 0 & 0 \end{pmatrix}$$

## Exercise Problems

Problems from the textbook: [E] 4.1 - 4.7

Exam problems: [Midterm 10/2018] Question 1-6

[Midterm 10/2022] Question 3,6,8

## Answers to Key Problems

### Problem 1.

- Eigenvalues  $\lambda_1 = -4$ ,  $\lambda_2 = 14$  and eigenvectors  $E_{-4} = \text{span}(\mathbf{v}_1)$  and  $E_{14} = \text{span}(\mathbf{v}_2)$ , where  $\mathbf{v}_1 = (-1,1)$  and  $\mathbf{v}_2 = (1,1)$ .
- Eigenvalues  $\lambda_1 = \lambda_2 = 3$  and eigenvectors  $E_3 = \text{span}(\mathbf{v}_1)$ , where  $\mathbf{v}_1 = (1,1)$ .
- No eigenvalues or eigenvectors.
- Eigenvalues  $\lambda_1 = \lambda_2 = 4$ ,  $\lambda_3 = 2$  and eigenvectors  $E_4 = \text{span}(\mathbf{v}_1, \mathbf{v}_2)$  and  $E_2 = \text{span}(\mathbf{v}_3)$ , where  $\mathbf{v}_1 = (0,1,0)$ ,  $\mathbf{v}_2 = (1,0,1)$ , and  $\mathbf{v}_3 = (-1,0,1)$ .
- Eigenvalues  $\lambda_1 = \lambda_2 = -1$ ,  $\lambda_3 = 2$  and eigenvectors  $E_{-1} = \text{span}(\mathbf{v}_1, \mathbf{v}_2)$  and  $E_2 = \text{span}(\mathbf{v}_3)$ , where  $\mathbf{v}_1 = (-1,1,0)$ ,  $\mathbf{v}_2 = (-1,0,1)$ , and  $\mathbf{v}_3 = (1,1,1)$ .
- Eigenvalues  $\lambda_1 = \lambda_2 = \lambda_3 = 0$  and eigenvectors  $E_0 = \text{span}(\mathbf{v}_1)$ , where  $\mathbf{v}_1 = (1,0,0)$ .

### Problem 2.

- Yes, with  $P = \begin{pmatrix} -1 & 1 \\ 1 & 1 \end{pmatrix}$ ,  $D = \begin{pmatrix} -4 & 0 \\ 0 & 14 \end{pmatrix}$
- No
- No
- Yes, with  $P = \begin{pmatrix} 0 & 1 & -1 \\ 1 & 0 & 0 \\ 0 & 1 & 1 \end{pmatrix}$ ,  $D = \begin{pmatrix} 4 & 0 & 0 \\ 0 & 4 & 0 \\ 0 & 0 & 2 \end{pmatrix}$
- Yes, with  $P = \begin{pmatrix} -1 & -1 & 1 \\ 1 & 0 & 1 \\ 0 & 1 & 1 \end{pmatrix}$ ,  $D = \begin{pmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 2 \end{pmatrix}$
- No

### Problem 3.

The eigenvalues of  $A$  are  $\lambda_1 = \lambda_2 = 5$ ,  $\lambda_3 = -1$  and  $\lambda_4 = -3$ .

### Problem 4.

- $A^m \rightarrow \begin{pmatrix} 1/5 & 1/5 \\ 4/5 & 4/5 \end{pmatrix}$  as  $m \rightarrow \infty$
- $A^m \rightarrow \begin{pmatrix} 2/3 & 2/3 \\ 1/3 & 1/3 \end{pmatrix}$  as  $m \rightarrow \infty$

### Problem 5.

- $-\lambda^3 + 6\lambda^2 - 4\lambda = 0$ ,  $\lambda = 0$  or  $\lambda = 3 \pm \sqrt{5}$
- $-\lambda^3 + 9\lambda^2 - 18\lambda + 8 = 0$ ,  $\lambda = 2$  or  $\lambda = (7 \pm \sqrt{33})/2$
- $-\lambda^3 = 0$ ,  $\lambda = 0$  (multiplicity 3)