

## Key Problems

### Oppgave 1.

Use Gaussian elimination to solve the linear systems with the following augmented matrices:

$$\text{a) } \left( \begin{array}{ccc|c} 1 & 3 & 4 & 11 \\ 2 & -1 & 3 & 3 \\ 3 & 2 & 5 & 12 \end{array} \right)$$

$$\text{b) } \left( \begin{array}{ccc|c} 1 & 3 & 4 & 11 \\ 2 & -1 & 3 & 3 \\ 3 & 2 & 7 & 12 \end{array} \right)$$

$$\text{c) } \left( \begin{array}{cccc|c} 1 & 1 & 1 & 1 & 8 \\ 1 & 3 & 1 & 5 & 28 \\ 2 & 4 & 2 & 9 & 48 \end{array} \right)$$

### Oppgave 2.

Determine how many solutions the linear system has:

$$\begin{aligned} x + y + 2z &= 6 \\ x + 2y + 4z &= 13 \\ x + 3y + 9z &= 24 \end{aligned}$$

Does the number of solutions change if we change the blue coefficient in the first equation? In that case, determine how the number of solutions changes with the blue coefficient.

### Oppgave 3.

We consider the homogeneous linear system with coefficient matrix

$$A = \begin{pmatrix} 1 & 1 & 4 & -1 \\ 5 & 5 & -1 & 4 \\ 7 & 6 & 3 & 3 \end{pmatrix}$$

Describe the set of solutions geometrically. How many degrees of freedom are there? Does this change if we change the red coefficient in the second row?

## Problems from the Workbook and Lecture Notes

Workbook [W]      1.1 - 1.18 (full solutions in the workbook)  
Lecture Notes [LN]    1.5 - 1.6, 1.9, 1.16 (solutions will be added soon)

## Answers to Key Problems

### Problem 1.

$$\text{a) } (x, y, z) = (1, 2, 1) \qquad \text{b) No solutions} \qquad \text{c) } (x, y, z, w) = (2 - z, 2, z, 4) \text{ with } z \text{ free}$$

### Problem 2.

There is one unique solution. The number of solutions only changes if the blue coefficient is  $-1$ , in which case there are no solutions. For any other value, there is a unique solution.

### Problem 3.

We have that  $\text{rk}(A) = 3$ , and there is  $n - \text{rk}(A) = 4 - 3 = 1$  degrees of freedom. Therefore the set of solutions is a straight line in  $\mathbb{R}^4$ . If we change the red coefficient, the rank of  $A$  remains  $\text{rk}(A) = 3$  unless the coefficient is  $6$ , in which case  $\text{rk}(A) = 2$ . Therefore, the set of solutions is a line for all values of the red coefficient except  $6$ , and in this case the set of solutions is a plane since the dimension is  $n - \text{rk}(A) = 4 - 2 = 2$ .